

CASE REPORT

Long-term Improvement after the Athens Protocol for Advanced Keratoconus with Significant Ectasia Progression in the Fellow Eye

¹Luis Claudio Pereira, ²Renata Siqueira, ³Bruno Valbon, ⁴Rodrigo Santos, ⁵Daniel Dawson
⁶Renato Ambrósio, ⁷Jorge Augusto Siqueira

ABSTRACT

To describe the long-term result of the Athens protocol (custom topography-guided advanced surface ablation followed by riboflavin-UVA collagen cross-linking in the same day) for progressive advanced keratoconus in the left eye, while the fellow right eye first presented with relatively mild keratoconus that was treated with a more conservative therapeutic approach of continuous contact lenses use with the goal for ectasia stabilization. Five years after the Athens protocol, a regression of over 10D (diopters) was observed on the sagittal curvature with significant regularization of corneal shape and improvement of distance corrected visual acuity (DCVA) in the left eye. The right eye, which was treated more conservatively, had an over 12D increase on sagittal curvature along with significant distortion of the corneal shape and reduction of DCVA.

Keywords: Keratoconus, Athens protocol, Contact lens.

How to cite this article: Pereira LC, Siqueira R, Valbon B, Santos R, Dawson D, Ambrósio R, Siqueira JA. Long-term Improvement after the Athens Protocol for Advanced Keratoconus with Significant Ectasia Progression in the Fellow Eye. *Int J Kerat Ect Cor Dis* 2014;3(1):40-43.

Source of support: Nil

Conflict of interest: None declared

INTRODUCTION

Keratoconus is a progressive, bilateral ectatic disorder in which the cornea becomes irregular and cone-shaped.¹ The ectatic condition is related to a progressive thinning and bulging due to chronic biomechanical failure of the corneal stroma. Corneal collagen cross-linking was developed in 1998 by Prof. Seiler and coworkers for augmenting corneal rigidity through the application of riboflavin (vitamin-B2) eye drops followed by ultraviolet A (UV-A) radiation.²

The application of customized topography-guided surface ablation has been reported in patients with stable or subclinical keratoconus.³⁻⁵ The Athens Protocol⁶ consists of the combination of same-day advanced custom surface ablation and riboflavin-UVA collagen cross-linking.⁷⁻⁹ The main goal of the procedure is to regularize the cornea,¹⁰ not to provide primary refractive correction, so that this approach is considered as therapeutic.¹¹

Special contact lens fitting, including rigid gas permeable (RGP) contact lenses, represents the main conservative therapeutic approach for visual rehabilitation of ectatic corneal diseases and is responsible for over 50% of cases.^{12,13} Contact lenses may provide the best possible visual acuity because it regulates the corneal surface along with the tear film.^{14,15} However, there is no evidence that contact lens stabilize or prevent ectasia progression, while there is some anecdotal evidence that it may be associated with a higher risk of ectasia progression due to continuous microtrauma from chronic RGP contact lens use.^{16,17}

CASE REPORT

A Caucasian 26-year-old presented with a history of decreased vision in his left eye since 2007. On examination, uncorrected distance visual acuity (UDVA) was 20/60 of the right eye and <20/400 of the left eye. Distance best corrected vision (DCVA) with spectacles was 20/20 in the right eye and 20/80 in the left eye. Fundus examination was within normal limits in both eyes. Goldmann applanation intraocular pressure (IOP) was 10 mm Hg in oculus dextrus (OD) and 7 mm Hg in oculus sinister (OS). Goldmann-correlated IOP (IOPg), corneal-compensated intraocular pressure (IOPcc), corneal hysteresis (CH), and corneal resistance factor (CRF) from the ocular response analyzer (ORA, Reichert, Buffalo, NY) were, respectively, 14.1, 11.4, 7.8 and 8.8 mm Hg of the right eye and 11.7, 7.6, 6.1 and 8.1 mm Hg of the left eye. Keratoconus was confirmed by placido disk-based corneal topography (iTrace) and rotating Scheimpflug corneal tomography (Pentacam, Oculus, Wetzlar, Germany). Wavefront optimized manifest refraction¹⁸ with data from iTrace (Tracey Technologies, Houston, TX) was $-1.25 - 2.75 \times 56^\circ = 20/20$ of the right eye and $-9.75 - 3.75 \times 1^\circ = 20/60$

^{1-4,7}Lecturer, ⁵Assistant Professor, ⁶Head

¹⁻⁶Department of Oftalmologia, Iora, Brazil

⁷Department of Refrativa e catarata, Iora: Instituto de Olhos Renato Ambrósio State, Rio de Janeiro, Brazil

Corresponding Author: Jorge Augusto Siqueira, Lecturer Department of Refrativa e catarata, Iora: Instituto de Olhos Renato Ambrósio State, Rio de Janeiro, Brazil, Phone: 552122644430, e-mail: jorgejass1@yahoo.com.br

of the left eye. The patient was fitted with RGP contact lens, providing a best corrected visual acuity of 20/15 of the right eye and 20/30 of the left eye.

Six months later, the patient returned complaining of decreased vision of the left, which failed to improve with contact lens wear. Wavefront (WF) optimized manifest refraction was $-1.50 - 2.75 \times 49^\circ = 20/20$ of the right eye, and $-11.75 - 6.00 \times 162^\circ = 20/80$ of the left. Significant progression of ectasia was demonstrated of the left eye with an increase of the maximal keratometric value (K_{max}) of 2.77 D, while the right eye remained stable.

Considering the progressive nature of the ectasia, the intolerance to further contact lens wear, the amount of irregularity, and the thinnest corneal thickness value of 456 μm of the left eye, the Athens Protocol⁶ was performed accordingly to the description by Kanellopoulos as an alternative for keratoplasty. Advanced surface ablation was performed with the Allegretto (Alcon-Wavelight, Erlangen, Germany). Epithelial removal was accomplished using the (phototherapeutic keratectomy) (PTK) mode set for 50 microns with a 6.5 mm optical zone and an 8.8 mm blend zone followed by subsequent same day custom topography-guided ablation (TCAT), based on oculyzer data, which removed 47 microns of tissue in the deepest ablation zone. Mitomycin C (0.005%) was applied for 45 seconds followed by copious irrigation with 50 cc of cold (4°C) BSS. Dextran-free 0.1% riboflavin solution was instilled every 2 minutes for 30 minutes. The yellow flare was verified before UVA irradiation at 3 mW/cm² for 30 minutes was completed. A bandage soft contact lens (SofLens 66, Bausch and Lomb Inc, Rocheste, New York) was placed until the cornea epithelialized on post-operative day 6 with no adverse event besides moderate pain in the first 3 days after the procedure. One month after the procedure, UCVA was 20/100. The patient had a second

opinion with another corneal subspecialist, who advised against corneal collagen cross-linking OD and who also indicated that continuous RGP contact lens wear in both eyes was necessary.

The patient returns in 2012 (4 years after the procedure) complaining of bad vision in both eyes and contact lens intolerance. UCVA was 20/100 of the right eye and 20/60 of the left eye. The wavefront (WF) optimized manifest refraction was $-13.50 - 7.00 \times 8^\circ = 20/70$ of the right eye and $-11.50 - 1.00 \times 169^\circ = 20/25$ of the left eye. Comparative subtraction front sagittal curvature maps from 2008 and 2012 are presented in Figure 1. Figures 2 and 3 include the elevation differential maps for OD and OS, respectively. 12.6D of corneal steepening is observed OD (Figs 1A and B) and 10.6D of flattening is observed OS (Figs 1C and D).

Figure 2 provides the front elevation maps for OD in 2008 and 2012. The same reference was chosen of 7.8 mm as best fit sphere (BFS) for the first evaluation. Over 80 microns of increase in elevation is observed as well as a decrease in central corneal thickness from 511 to 455 μm (at pupil center) and an increase in ACD (anterior chamber depth) (ACD) from 3.54 to 3.87 mm.

Figure 3 provides the front elevation maps for OS in 2008 and 2012. The same reference was chosen of 7.04 mm as best fit sphere (BFS) for the first evaluation. 36 microns of decrease in elevation was observed and a decrease in central keratometric astigmatism from 5.9D to 0.3D. A mild decrease in central corneal thickness from 478 to 443 μm (at pupil center) and an decrease in anterior chamber depth (ACD) from 3.87 to 3.77 mm.

The patient was advised about the need for therapeutic surgery for OD and surgical options, including the possibility of keratoplasty,¹⁹ and phakic intraocular lens (IOL) as a refractive option for OS.




Fig. 1: Sagittal curvature differential maps




Fig. 2: Right eye: Maps of front elevation (A: Postoperative; B: Preoperative) and the difference A-B, with the same BFS




Fig. 3: Left eye: Maps of front elevation (A: Postoperative; B: Preoperative) and the difference A-B, with the same BFS

DISCUSSION

The patient presented with advanced progressive keratoconus of the left eye, which was treated by the Athens protocol, and the right eye had a mild and relatively stable keratoconus presentation with good distance corrected visual acuity with spectacles. The patient had a second opinion, who advised against cross-linking of the right eye and to continuously wear RGP contact lens OD. This case illustrates that the use of contact lens does not stop the progression of keratoconus. Moreover, continuous microtrauma related to continuous contact lens wear may be an important factor to accelerate ectasia progression in some patients.^{16,17} After 4 years, a regression of more than 10D of astigmatism of the left eye was associated with regularization of the corneal surface and improvement in the BCVA after the Athens protocol. This result is consistent with the literature.^{6,20-22}

CONCLUSION

The Athens protocol is a therapeutic alternative for patients with progressive keratoconus to prevent or even delay keratoplasty. This is not a refractive procedure. Refractive correction with a phakic IOL becomes a possibility, if best spectacle corrected vision is good. However, the progressive nature of ectatic corneal diseases should be considered and patient education is fundamental. Surgery for keratoconus should be indicated when visual rehabilitation is not possible with glasses or contact lenses and when there is documented or a significant risk of ectasia progression. While contact lenses provide the best possible treatment for the irregular astigmatism, there is no evidence to support the benefit of contact lens for decreasing the risk of ectasia progression. In fact, contact lens fitting should be considered with caution, if spectacles provide adequate visual acuity.

REFERENCES

1. Rabinowitz YS. Keratoconus. *Surv Ophthalmol* 1998;42(4):297-319.
2. Spoerl E, Huhle M, Seiler T. Induction of cross-links in corneal tissue. *Exp Eye Res* 1998;66(1):97-103.
3. Alpíns N, Stamatelatos G. Customized photoastigmatic refractive keratectomy using combined topographic and refractive data for myopia and astigmatism in eyes with forme fruste and mild keratoconus. *J Cataract Refract Surg* 2007;33(4):591-602.
4. Koller T, Iseli HP, Donitzky C, Ing D, Papadopoulos N, Seiler T. Topography-guided surface ablation for forme fruste keratoconus. *Ophthalmology* 2006;113(12):2198-2202.
5. Cennamo G, Intravaja A, Boccuzzi D, Marotta G. Treatment of keratoconus by topography-guided customized photorefractive keratectomy: two-year follow-up study. *J Refract Surg* 2008;24(2):145-149.
6. Kanellopoulos AJ. Comparison of sequential vs same-day simultaneous collagen cross-linking and topography-guided PRK for treatment of keratoconus. *J Refract Surg* 2009;25(9):S812-818.
7. Wasilewski D, Mello GH, Moreira H. Impact of Collagen Cross-linking on Corneal Sensitivity in Keratoconus Patients. *Cornea* 2013;32(7):899-902.
8. Viswanathan D, Males J. Prospective longitudinal study of corneal collagen cross-linking in progressive keratoconus. *Clin Experiment Ophthalmol* 2013;41(6):531-536.
9. Cheema AS, Mozayan A, Channa P. Corneal collagen cross-linking in refractive surgery. *Curr Opin Ophthalmol* 2012;23(4):251-256.
10. Oshika T, Tomidokoro A, Maruo K, Tokunaga T, Miyata N. Quantitative evaluation of irregular astigmatism by fourier series harmonic analysis of videokeratography data. *Invest Ophthalmol Vis Sci* 1998;39(5):705-709.
11. Ambrósio Jr. R. Cirurgia refrativa terapêutica: por que diferenciar? *Revista Brasileira de Oftalmologia* 2013;72(4):85-86.
12. Wagner H, Barr JT, Zadnik K. Collaborative Longitudinal Evaluation of Keratoconus (CLEK) Study: methods and findings to date. *Cont Lens Anterior Eye* 2007;30(4):223-232.
13. Weed KH, Macewen CJ, McGhee CN. The Dundee University Scottish Keratoconus Study II: a prospective study of optical and surgical correction. *Ophthalmic Physiol Opt* 2007;27(6):561-567.
14. Mandathara PS, Fatima M, Taureen S, Dumpati S, Ali MH, Rathi V. RGP contact lens fitting in keratoconus using FITSCAN technology. *Cont Lens Anterior Eye* 2013;36(3):126-129.
15. Fatima T, Acharya MC, Mathur U, Barua P. Demographic profile and visual rehabilitation of patients with keratoconus attending contact lens clinic at a tertiary eye care centre. *Cont Lens Anterior Eye* 2010;33(1):19-22.
16. McMonnies CW. Mechanisms of rubbing-related corneal trauma in keratoconus. *Cornea* 2009;28(6):607-615.
17. Barr JT, Zadnik K, Wilson BS, Edrington TB, Everett DF, Fink BA, Shovlin JP, Weissman BA, Seigmund K, Gordon MO. Factors associated with corneal scarring in the Collaborative Longitudinal Evaluation of Keratoconus (CLEK) Study. *Cornea* 2000;19(4):501-507.
18. Ambrósio Junior R, Caldas DL, Silva RSd, Pimentel LN, Valbon BdF. Impacto da análise do "wavefront" na refratometria de pacientes com ceratocone. *Revista Brasileira de Oftalmologia* 2010;69:294-300.
19. Vazirani J, Basu S. Keratoconus: current perspectives. *Clin Ophthalmol* 2013;7:2019-2030.
20. Kremer I, Aizenman I, Lichter H, Shayer S, Levinger S. Simultaneous wavefront-guided photorefractive keratectomy and corneal collagen cross-linking after intrastromal corneal ring segment implantation for keratoconus. *J Cataract Refract Surg* 2012;38(10):1802-1807.
21. Spadea L. Collagen cross-linking for ectasia following PRK performed in excimer laser-assisted keratoplasty for keratoconus. *Eur J Ophthalmol* 2012;22(2):274-277.
22. Kymionis GD, Grentzelos MA, Portaliou DM, Karavitaki AE, Krasia MS, Dranidis GK, Kozobolis VP. Photorefractive keratectomy followed by same-day corneal collagen cross-linking after intrastromal corneal ring segment implantation for pellucid marginal degeneration. *J Cataract Refract Surg* 2010;36(10):1783-1785.